

Physics of Complex Magnetic Materials: Quasiparticle Many-Body Dynamics

A. L. Kuzemsky *

*Bogoliubov Laboratory of Theoretical Physics,
Joint Institute for Nuclear Research,
141980 Dubna, Moscow Region, Russia.*

Abstract

A brief survey of the author's works on the quantum theory of magnetism. Theoretical foundation and applications of the generalized spin-fermion (sp-d) exchange lattice model to various magnetic systems, e.g., rare-earth metals and compounds, and magnetic and diluted magnetic semiconductors are discussed briefly. The main emphasis is put on the dynamic behavior of two interacting subsystems, the localized spins and spin density of itinerant carriers. A nonperturbative many-body approach was used to describe the quasiparticle dynamics. The approach permits to investigate and clarify the role of various interactions and disorder effects in combined quantum models of magnetism in unified and coherent fashion.

The existence and properties of localized and itinerant magnetism in metals, oxides, and alloys, and their interplay is an interesting but not yet fully understood problem of quantum theory of magnetism[1, 2]. The behavior and the true nature of the electronic and spin states, and their quasiparticle dynamics are of central importance to the understanding of physics of correlated systems such as magnetism and Mott-Hubbard metal-insulator transition in metals and oxides, magnetism and heavy fermions in rare-earths compounds, and anomalous transport properties in perovskite manganites. This class of systems is characterized by complex, many-branch spectra of elementary excitations. Moreover, the correlation effects (competition and interplay of Coulomb correlation, direct or indirect exchange, sp-d hybridization, electron-phonon interaction, disorder, etc.) are essential[3, 4]. These materials are systems of great interest both intrinsically and as a possible source of understanding the magnetism of matter generally [5, 6]. There has been considerable interest in identifying the microscopic origin of quasiparticle states [3, 4, 6] in these systems and a few model approaches have been proposed. Many magnetic and electronic properties of rare-earth metals and compounds, and magnetic semiconductors [7] and related complex magnetic materials may reasonably be interpreted

*E-mail:kuzemsky@theor.jinr.ru; <http://theor.jinr.ru/~kuzemsky>

in terms of combined spin-fermion models which include interacting spin and charge subsystems [8, 9, 10]. This approach permits one to describe significant and interesting physics, e.g., bound states and magnetic polarons [9], anomalous transport properties, etc.

The problem of adequate physical description within various types of spin-fermion model has intensively been studied during the last decades, especially in the context of magnetic and transport properties of rare-earth and transition metals and their compounds and magnetic semiconductors.

Substances which we refer to as magnetic semiconductors, occupy an intermediate position between magnetic metals and magnetic dielectrics. Magnetic semiconductors are characterized by the existence of two well defined subsystems, the system of magnetic moments which are localized at lattice sites, and a band of itinerant or conduction carriers (conduction electrons or holes). Typical examples are the Eu-chalcogenides, where the local moments arise from 4f electrons of the Eu ion, and the spinell chalcogenides containing Cr^{3+} as a magnetic ion [7]. There is experimental evidence of a substantial mutual influence of spin and charge subsystems in these compounds. This is possible due to the $sp - d(f)$ exchange interaction of the localized spins and itinerant charge carriers. More recent efforts have been directed to the study of the properties of diluted magnetic semiconductors (DMS)[10]. Further attempts have been made to study and exploit carriers which are exchange-coupled to the localized spins. The effect of carriers on the magnetic ordering temperature is found to be very strong in DMS. Diluted magnetic semiconductors are mixed crystals in which magnetic ions (usually Mn^{++}) are incorporated in a substitutional position of the host (typically a II-VI or III-V) crystal lattice. The diluted magnetic semiconductors offer a unique possibility for a gradual change of the magnitude and sign of exchange interaction by means of technological control of carrier concentration and band parameters. This field is very active and there are many aspects to the problem. A lot of materials were synthesized and tested. The new material design approach to fabrication of new functional diluted magnetic semiconductors resulted in producing a variety of compounds. The presence of the spin degree of freedom in DMS may lead to a new semiconductor spin electronics which will combine the advantages of the semiconducting devices with the new features due to the possibilities of controlling the magnetic state. However, the coexistence of ferromagnetism and semiconducting properties in these compounds require a suitable theoretical model which would describe well both the magnetic cooperative behavior and the semiconducting properties as well as a rich field of interplay between them. The majority of theoretical papers on DMS studied their properties mainly within the mean field approximation and continuous media terms. In a picture like this the disorder effects, which play an essential role, can be taken into account roughly only. Moreover, there are different opinions on the intrinsic origin and the nature of disorder in DMS. Thus, many experimental and theoretical investigations call for a better understanding of the relevant physics and the nature of solutions (especially magnetic) within the lattice spin-fermion model[8, 9, 10]. For treating the problems we used a nonperturbative many-body approach, the formalism of the method of Irreducible Green Functions (IGF) [4]. This IGF method allows one to

describe quasiparticle spectra with damping for many-particle systems on a lattice with complex spectra and a strong correlation in a very general and natural way. This scheme differs from the traditional method of decoupling of an infinite chain of equations [1] and permits a construction of the relevant dynamic solutions in a self-consistent way at the level of the Dyson equation without decoupling the chain of equations of motion for the GFs.

In paper [9] the concepts of bound and scattering states were analysed and developed to elucidate the nature of itinerant charge carrier states in magnetic semiconductors and similar complex magnetic materials. By contrasting the scattering and bound states of carriers within the $s - d$ exchange model, the nature of bound states at finite temperatures was clarified. The free magnetic polaron at certain conditions is realized as a bound state of the carrier (electron or hole) with the spin wave. Quite generally, a self-consistent theory of a magnetic polaron was formulated within the IGF method which was used to describe the quasiparticle many-body dynamics at finite temperatures. Within the above approach we elaborated a self-consistent picture of dynamic behavior of two interacting subsystems, the localized spins and the itinerant charge carriers. In particular, it was shown that the relevant generalized mean fields emerges naturally within our formalism. At the same time, the correct separation of elastic scattering corrections permitted one to consider the damping effects (inelastic scattering corrections) self-consistently. The damping of magnetic polaron state, which is quite different from the damping of the scattering states, finds a natural interpretation within the present self-consistent scheme.

In paper [10], we applied the IGF formalism to consider quasiparticle spectra for the lattice spin-fermion model consisting of two interacting subsystems. It was the purpose of that paper to explore more fully the notion of Generalized Mean Fields (GMF) [4] which may arise in the system of interacting localized spins (including effects of disorder) and lattice fermions to justify and understand the nature of the relevant mean fields. Background and applications of the generalized spin-fermion (sp-d) exchange model to magnetic and diluted magnetic semiconductors were discussed in some detail. The capabilities of the model to describe quasiparticle spectra were investigated. The key problem of most of the work was the formation of spin excitation spectra under various conditions on the parameters of the model. In paper [10], we concentrated on the description of the magnetic excitation spectra and treated the disorder effects in the simplest VCA (virtual crystal approximation) to emphasize the need for a suitable definition of the relevant generalized mean fields and for internal self-consistency in the description of the spin quasiparticle many-body dynamics.

Thus, we were able to calculate the quasiparticle spectra and GMF of the magnetic semiconductors consisting of two interacting charge and spin subsystem within the lattice spin-fermion model to analyze the role and influence of the Coulomb correlation, exchange, and effects of disorder in a unified and coherent fashion.

References

- [1] S. V. Tyablikov, *Methods in the Quantum Theory of Magnetism*, Plenum Press (New York, 1967).
- [2] A. L. Kuzemsky, Fundamental Principles of the Physics of Magnetism and the Problem of Localized and Itinerant Electronic States, Communication JINR, E17-2000-32, Dubna, 2000.
- [3] A. L. Kuzemsky, Quasiparticle Many-Body Dynamics of Highly Correlated Electronic Systems, Molecular Physics Reports, **17**, 221 (1997). (cond-mat/9704028)
- [4] A. L. Kuzemsky, Irreducible Green Functions Method and Many Particle Interacting Systems on a Lattice, Rivista Nuovo Cimento **25**, 1 (2002). (cond-mat/0208219)
- [5] A. L. Kuzemsky, Quantum Protectorate and Microscopic Models of Magnetism, Intern. J. Modern Phys. B **16**, 803 (2002). (cond-mat/0208277)
- [6] A. L. Kuzemsky, Itinerant Antiferromagnetism of Correlated Lattice Fermions, Physica A **267**, 131 (1999). (cond-mat/0208242)
- [7] E. L. Nagaev, *Physics of Magnetic Semiconductors*, MIR Publ., (Moscow, 1985).
- [8] A. L. Kuzemsky, Spectral Properties of Generalized Spin-Fermion Models, Intern. J. Modern Phys. B **13**, 2573 (1999). (cond-mat/0208277)
- [9] A. L. Kuzemsky, Bound and Scattering States of Itinerant Charge Carriers in Complex Magnetic Materials, Intern. J. Modern Phys. B **18**, 3227 (2004). (cond-mat/0408404)
- [10] A. L. Kuzemsky, Role of Correlation and Exchange for Quasiparticle Spectra of Magnetic and Diluted Magnetic Semiconductors, Physica B **355**, 318 (2005).